

## Influenza Surveillance 2015–2016

Week 7 (Feb.14–20, 2016)

Data extracted Feb. 26, 2016 at 11:00 am

### Laboratory-confirmed influenza cases this week:

- **50** cases of influenza A ↑
- **2** case of influenza B ↑
- **13.0%** positivity ↑

### Since Sept. 1, 2015:

- **137** cases of influenza A
- **11** case of influenza B

### Laboratory

### Cases and cumulative incidence rates (cases per 100,000 population) since Sept. 1, 2015:

- **Winnipeg: 45 (6.0)**
- **Southern: 21 (11.0)**
- **Interlake-Eastern: 14 (11.1)**
- **Prairie Mountain: 33 (19.7)**
- **Northern: 35 (46.5)**

### Regional Health Authority

### Severe outcomes associated with laboratory-confirmed influenza diagnosis in this week:

- **4** hospitalizations →
- **1** ICU admission →
- **0** deaths →

### Since Sept. 1, 2015:

- **18** hospitalizations
- **5** ICU admissions
- **0** deaths

### Severity

### Calls to Influenza Service at Health Links–Info Santé this week: **16** ↑

### Percent of visits to sentinel physicians due to ILI this week: **1.4%** ↑

### In Emergency Department this week:

- **190** ILI cases per day on average ↑
- **19–27%** of visits due to ILI ↑

### Influenza-Like-Illness (ILI)

### Laboratory-confirmed outbreaks this week:

- **0** outbreak of influenza A ↓
- **0** outbreaks of influenza B →

### Since Sept. 1, 2015:

- **3** outbreaks of influenza A
- **0** outbreaks of influenza B

### Outbreak

### Oseltamivir dispensed from community retail pharmacies:

- This week: **84** units ↑
- Since Oct. 1, 2015: **508** units

### Isolates resistant to antiviral since Sept. 1, 2015 in Manitoba:

- **Oseltamivir: 1**
- **Zanamivir: 0**

### Antiviral

### As of Jan. 31, 2016:

- **Percent of Manitobans immunized with the seasonal influenza vaccine: 22%**

### Please note:

- **FluMist® Quadrivalent (AstraZeneca)** vaccines expired and are not available any more this season.
- **Fluzone® Quadrivalent (Sanofi Pasteur)** vaccines are still available.

### Immunization

This week, influenza activity in Manitoba continued to increase and the largest increase in influenza cases occurred in Prairie Mountain. ILI activity in Emergency Departments increased significantly from last week, especially among children below 10 years of age and seniors above 65. Incidence rate was much higher among young children below 5. They accounted for one third of all hospitalized influenza patients and seniors above 65 accounted for another third.

Nationally, more laboratory detections and outbreaks of influenza were reported. Young and middle age adults accounted for an increasing proportion of hospitalizations. In British Columbia, Alberta and Ontario, the increase in laboratory detections of respiratory viruses including influenza seemed to slow down compared to two weeks ago. In those provinces, influenza accounted for 70% of all respiratory viruses detected.


Influenza activity in the United States also increased this week and one influenza-associated pediatric death was reported. On February 25, 2016, the WHO announced its recommended components for the 2016–2017 northern hemisphere influenza vaccine.

### Summary: Increasing activity


**Note.** Numbers are subject to change. Missed events in the current weekly report due to a delay of submission to MHLS will be reported in later weeks when data become available.

### Laboratory Surveillance

Reports of influenza nucleic acid detection, culture isolation, and enzyme immunoassay (EIA) detections are received from Cadham Provincial Laboratory (CPL) and occasionally other laboratories. These reports are forwarded to Epidemiology and Surveillance (E&S) within 24 hours of confirmation. CPL also performs testing for other respiratory viruses including parainfluenza, RSV, adenovirus, rhinovirus, coronavirus, enterovirus, and bocavirus, which are reported to E&S on a weekly basis.



**Figure 1. Laboratory-confirmed influenza A cases by week, Manitoba, 2015–2016**



**Figure 2. Laboratory-confirmed influenza B cases by week, Manitoba, 2015–2016**




Figure 3. Influenza A and B cases by age group, Manitoba, 2015–2016

### Health Links – Info Santé

Health Links–Info Santé is a 24-hour, 7-days a week telephone information service. It is staffed by registered nurses with the knowledge to provide answers to health care questions and guidance to appropriate care over the phone. When a caller phones Health Links–Info Santé and selects Influenza Service, they are given an option to select information on (1) the groups of individuals who are at an increased risk of serious illness, (2) how to arrange an influenza vaccine, (3) the annual influenza immunization campaign, or (4) the management of influenza and its potential complications.

The number of calls to Health Links – Info Santé continued to be low. Overall, there were less calls in 2015–2016 than in previous seasons.





Figure 4. Calls to Health Links – Info Santé, Manitoba, 2015–2016

### ILI

#### ILI visits to sentinel physicians

Manitoba participates in *FluWatch*, the Canada’s national surveillance system co-ordinated by Public Health Agency of Canada (PHAC), which monitors the spread of influenza and ILI on a year-round basis. *FluWatch* consists of a network of laboratories, hospitals, doctor’s offices and provincial and territorial ministries of health. In 2015–2016, there were 19 sentinel physicians recruited throughout Manitoba. They are requested to report to *FluWatch* weekly. E&S receives weekly reports from *FluWatch* which present the ILI rate for Manitoba and for each of the participating sentinel physicians. Note that the reporting sentinel physicians are different by week and their reports may not be representative of ILI activity across the province.

The weekly percentage of patient visits to sentinel physicians due to ILI this season continued to be low.




**Figure 5 Percent of patient visits to sentinel physicians due to ILI by week, Manitoba, 2015–2016**

**ILI visits to Emergency Rooms**

E&S receives the summary report for the daily ILI related visits to Emergency Department at Winnipeg Regional Health Authority (WRHA) on a weekly basis. ILI cases are defined as patients whose triage chief complaints contain either of these symptoms: weakness, shortness of breath, cough, headache, fever, cardiac/respiratory arrest, sore throat, and upper respiratory tract infection complaints.

The number of ILI cases and as % of total visits in Emergency Department increased significantly this week compared to the previous week.



**Figure 6 ILI cases as % of all visits in Emergency Department at WRHA, Manitoba**

### Antiviral dispensing

Daily units of antiviral drug, Oseltamivir (Tamiflu®), dispensed to Manitoba residents during an influenza season are reported to E&S from Drug Programs Information Network (DPIN) on a weekly basis since October 1 each season. Only drugs dispensed from community retail pharmacies are included in this report. Antiviral drugs dispensed to in-patients or through nursing stations could not be included due to lack of data.




Figure 7 Units of Oseltamivir dispensed by week, Manitoba, 2015–2016

### Antiviral Resistance

Influenza and Respiratory Viruses Section of National Microbiology Laboratory (NML) undertakes enhanced surveillance, investigations, and research on influenza and other respiratory pathogens. A random sample of positive influenza specimens isolated by culture is referred from each provincial laboratory to NML for strain characterization and antiviral resistance testing. The aggregate level information is then shared with provinces and territories on a weekly basis.

To date, all influenza viruses tested except one were sensitive to Oseltamivir and all were sensitive to Zanamivir. In comparison, almost all viruses tested were resistant to Amantadine.

Table 1. Antiviral resistance of isolates by influenza type and subtype since September 1, 2015 in Canada and Manitoba, 2015–2016

		Oseltamivir		Zanamivir		Amantadine	
		# Resistant	# Sensitive	# Resistant	# Sensitive	# Resistant	# Sensitive
<b>Canada</b>	A(H3N2)	0	116	0	116	125	1
	A(H1N1)	1	229	0	130	210	0
	B	0	119	0	119	N/A	N/A
<b>Manitoba</b>	A(H3N2)	0	0	0	0	0	0
	A(H1N1)	0	5	0	5	5	0
	B	0	2	0	2	N/A	N/A

N/A = Not applicable

### Immunization

As per World Health Organization (WHO), all seasonal quadrivalent influenza vaccines for 2015–2016 in the northern hemisphere contain:

- A/Switzerland/9715293/2013(H3N2)-like virus;

- A/California/7/2009(H1N1)pdm09-like virus;
- B/Phuket/3073/2013-like virus.
- B/Brisbane/60/2008-like virus.

For the 2015–2016 influenza season, MHLS is allotted quadrivalent inactivated vaccine (QIV), Fluzone® Quadrivalent (Sanofi Pasteur), and quadrivalent live attenuated influenza vaccine (QLAIV) FluMist® Quadrivalent (AstraZeneca), as part of the province’s Publicly-Funded Seasonal Influenza Immunization Program.

## Circulating Strain

NML antigenically characterizes influenza viruses received from Canadian laboratories year-round. In Manitoba, a random sample of positive influenza specimens isolated by culture is referred from CPL to NML.

Since September 1, 2015, NML has characterized 521 influenza A and B viruses.

- 122 influenza A(H3N2) viruses:
  - 27 influenza A(H3N2) viruses were antigenically characterized as A/Switzerland/9715293/2013, the influenza A(H3N2) component in the 2015–2016 influenza vaccine.
  - 95 influenza A(H3N2) viruses did not grow to sufficient hemagglutination titers for antigenic characterization by hemagglutination inhibition assays. Therefore, genetic characterization was performed. Sequence analyses showed that those influenza A(H3N2) viruses belonged to a genetic group in which most viruses were antigenically related to A/Switzerland/9715293/2013.
- 268 influenza A(H1N1) viruses:
  - 268 influenza A(H1N1) viruses characterized were antigenically similar to A/California/7/2009, the influenza A(H1N1) component in the vaccine.
- 131 influenza B viruses:
  - 46 influenza B viruses characterized were antigenically similar to B/Phuket/3073/2013 (Yamagata lineage), the influenza B component in the vaccine.
  - 85 influenza B viruses were characterized as B/Brisbane/60/2008-like (Victoria lineage), the influenza B component in the quadrivalent vaccine.

**Table 2 Influenza Strain Characterization reported by NML since September 1, 2015, Canada, 2015–2016**

Strain	Number of viruses	
	Canada	Manitoba
A/Switzerland/9715293/2013(H3N2)-like	27	0
A/California/7/2009(H1N1)-like	268	5
B/Phuket/3073/2013-like	46	1
B/Brisbane/60/2008-like	85	2

## Abbreviations

**CPL = Cadham Provincial Laboratory**  
**E&S = Epidemiology and Surveillance**  
**ICU = Intensive Care Unit**  
**ILI = Influenza Like Illness**  
**LTCF = Long Term Care Facility**  
**MHLS = Manitoba Health, Healthy Living and Seniors**  
**NML = National Microbiology Laboratory**  
**PHAC = Public Health Agency of Canada**  
**RHA = Regional Health Authority**  
**WHO = World Health Organization**  
**WRHA = Winnipeg Regional Health Authority**

## Explanatory Notes and Definitions

### Cumulative data

Cumulative data include updates to previous weeks. Due to reporting delays or amendments, the sum of weekly report totals may not add up to cumulative totals.

### Data extraction date

Manitoba-specific information contained within this update is based on data confirmed in SIS databases at 11:00 am on the date of data extraction.

### Epidemiology week

Time trends in this report were analyzed by [epidemiology week](#), a schedule used by the national FluWatch program coordinated by the Public Health Agency of Canada (PHAC).

### Incidence rate

Incidence rate measures the frequency with which influenza occurs in a region. It is calculated as the total number of new cases this influenza season multiplied by 10,000 and divided by the total population in each region. Regional populations as of June 1, 2014 are provided by Information Management & Analytics at MHLS.

### ILI in the general population

Acute onset of respiratory illness with fever and cough and with one or more of the following – sore throat, arthralgia, myalgia, or prostration, which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

### ILI outbreaks

Schools: Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI.

Hospitals and residential institutions: Two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case.

Other settings: Two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case; i.e. workplace, closed communities.

### Specimen collection date

The date the laboratory specimen was taken is used to assign cases to the appropriate week in this report. However, hospitalized/ICU cases are reported based on laboratory report date.

**Epidemiology and Surveillance**  
**Manitoba Health, Healthy Living and Seniors**  
[flusurveillance@gov.mb.ca](mailto:flusurveillance@gov.mb.ca) / (204)786-7335

For other Epidemiology and Surveillance reports, please view the Manitoba Health internet website:  
<http://www.gov.mb.ca/health/publichealth/surveillance/index.html>

For national surveillance data, refer to:  
<http://www.phac-aspc.gc.ca/fluwatch/index-eng.php>